

SPECIFICATION SHEET

VERSION 4 **VERSION DATE** 19/09/22 **PRINT DATE** 22/08/25

PRODUCT ACETYL METHYL CARBINOL

APPLICATIONS Industrial raw material not intended for direct use.

IUPAC NAME 3-HYDROXI-2-BUTANONE

SYNONYMS 3-HYDROXY-2-BUTANONE, ACETOIN, ACETYLMETHYLCARBINO

FORMULA C₄ H₈ O₂ **MOLECULAR WEIGHT** 88 **CAS** 513-86-0 208-174-1 **EINECS** 2008 **FEMA** 749 CoF FDΔ 182 60 TARIFF NUMBER 2914409090

FOOD GRADE CERTIFICATE AVAILABLE ON REQUEST

SPECIFICATIONS

VERSION 4 VERSION DATE 19/09/22

ASPECT AT 20°C CLEAR MOBILE LIQUID TO SOLID (DIMER)

COLOUR COLORLESS TO PALE YELLOW

 ODOUR
 BUTTER-LIKE

 TASTE
 BUTTER-LIKE

 DENSITY AT 20°C (g / ml)
 0,998-1,022

 DENSITY 25/25
 0,995-1,019

 REFRACTIVE INDEX nD20
 1,4170-1,4200

REFRACTIVE INDEX nD25 OPTICAL ROTATION (°) /
BOILING POINT (°C) 143

MELTING POINT (°C) CA. 15 (MONOMER);110 (DIMER)

 FLASH POINT (°C)
 50

 ASSAY (% GC)
 > 97

 ACID VALUE (mg KOH/g)
 < 2</td>

SOLUBILITYSOLUBLE IN WATERVAPOUR PRESSURENOT DETERMINEDOTHER SOLUBILITIESSOLUBLE IN ETHANOL

OTHER TECHNICAL DATA PURGE HEADSPACE WITH NITROGEN. RECOMMENDED NOT TO PACK IN

EPOXY-COATED STEEL DRUMS.

STABILITY AND STORAGE

KEEP IN TIGHTLY CLOSED CONTAINER IN A COOL AND DRY PLACE, PROTECTED FROM LIGHT. WHEN STORED FOR MORE THAN 24 MONTHS, QUALITY SHOULD BE CHECKED BEFORE USE.

THE PRODUCT TENDS TO DIMERIZE UPON STORAGE; THIS PROCESS CAN BE REVERSED BY HEATING NOT ABOVE 60°C

WE BELIEVE THE ABOVE INFORMATION TO BE CORRECT BUT WE DO NOT PRESENT IT AS ALL INCLUSIVE AND AS SUCH SHOULD BE USED AS A GUIDE. IT IS THE SOLE RESPONSIBILITY OF THE RECEIVER OF OUR PRODUCT TO COMPLY WITH EXISTING REGULATIONS AND INSTRUCTIONS. THIS PRODUCT HAS NOT BEEN FILTERED DUE TO TECHNICAL REASONS. WE RECOMMEND NOT TO USE IT UNLESS FILTERED WHEN USED AS A RAW MATERIAL IN FOOD INTENDED FOR FINAL CONSUMERS.

PRODASYNTH SAS